

Appendix A

**List of fishes collected by the Meisyo Maru #128 in 1993,
expressed in relative abundance in weight by seamount depth zone,
with related information**

Seamount surveyed Depth zone surveyed	C-H 200–400m	Colahan 200–400m	Kammu 200–400m	Yuryaku 400–700m	Yuryaku 200–400m	Koko 200–400m	Total
Operation start date (UTC) for each strata	1993/11/19	1993/11/16	1993/11/10	1993/11/7	1993/11/7	1993/11/19	
Type of trawl, bottom or mid-water.	Bottom	Bottom	Bottom	Bottom	Bottom	Bottom	
Type of trawl, single, double or	Single	Single	Single	Single	Single	Single	
Typical height of net opening (m).	5	5	5	5	5	5	
Typical width of net opening (m).	25	25	25	25	25	25	
Mesh size of the cod-end net (mm)	4	4	4	4	4	4	
Mesh type (diamond, square, etc).	Diamond	Diamond	Diamond	Diamond	Diamond	Diamond	
Number of effective trawl operation	1	8	23	3	5	16	
Total duration of tows (min)	4	77	238	30	48	248	
Total estimated duration of seabed contact (min)	4	77	238	30	48	248	
Total estimated area of seabed contact (km ²)	0.0093	0.1783	0.5510	0.0695	0.1111	0.5741	
TOTAL CATCH BY WEIGHT (kg)	432.5	3644.5	6736.1	293.8	12180.3	5616.6	28903.7
<i>Etmopterus pusillus</i>	15.3	7.8	308.4	104.1	14.4	40.9	490.8
<i>Squalus mitsukurii</i>	0.0	0.0	94.1	0.0	0.0	58.7	152.8
<i>Chimaera spp.</i>	0.0	0.0	0.0	0.0	0.0	13.9	13.9
<i>Meadia abyssalis</i>	0.6	3.2	14.7	0.5	20.4	0.5	39.9
<i>Chlorophthalmus spp.</i>	0.0	0.1	18.7	3.6	0.0	0.2	22.6
<i>Argyropelecus aculeatus</i>	0.0	0.0	0.3	0.0	0.0	0.0	0.3
<i>Myctophidae sp.</i>	0.0	0.0	0.0	0.0	0.0	0.0	0.0
<i>Polymixia japonica</i>	0.0	0.8	14.6	5.5	0.2	16.1	37.1
<i>Physiculus spp.</i>	0.6	8.1	197.2	10.8	5.2	1347.0	1568.9
<i>Macrouridae sp.</i>	0.1	0.1	12.9	2.8	0.2	1.5	17.5
<i>Lophiomus micanthus</i>	0.1	4.2	73.8	14.3	24.4	83.5	200.2
<i>Chaunax spp.</i>	0.0	0.2	18.1	2.7	20.3	38.6	79.7
<i>Malthopsis spp.</i>	0.0	0.0	0.0	0.0	0.0	0.3	0.4
<i>Bervx splendens</i>	24.5	1275.9	1242.8	50.2	11591.2	16.9	14201.5
<i>Beryx decadactylus</i>	0.0	0.0	3.3	0.0	0.0	48.9	52.2
<i>Scorpaena spp.</i>	0.0	0.0	0.1	0.5	0.0	0.0	0.6
<i>Satyrichthys engypterus</i>	0.0	0.0	2.6	0.6	0.6	0.2	3.9
<i>Macroramphosus spp.</i>	0.0	2.8	6.5	0.0	0.0	143.2	152.4
<i>Zenopsis nebulosa</i>	15.7	284.9	3933.3	1.9	15.7	845.1	5096.6
<i>Zenion japonicum</i>	0.0	0.0	4.5	0.0	0.0	0.0	4.5
<i>Parazen pacificus</i>	0.0	0.0	0.5	0.0	0.0	0.0	0.5
<i>Hoplostethus crassispinus</i>	0.7	0.1	2.3	7.7	0.1	0.0	11.0
<i>Congriscus megastomus</i>	0.7	0.2	1.6	0.1	0.5	0.3	3.3
<i>Helicolenus spp.</i>	0.0	0.0	141.1	0.2	0.1	523.3	664.7
<i>Epigonus denticulatus</i>	11.6	1.3	90.9	64.8	5.1	2327.9	2501.5
<i>Plectranthias kelloggi</i>	0.0	1.0	7.7	0.0	0.0	2.6	11.3
<i>Sympodus maunaloa</i>	0.1	1.5	4.4	0.0	0.0	0.5	6.4
<i>Decapterus tabl</i>	0.0	0.0	0.0	0.0	0.0	0.6	0.6
<i>Emmelichthys struhsakeri</i>	0.0	0.1	0.0	0.0	0.0	0.0	0.1
<i>Cookeolus japonicus</i>	0.0	0.3	0.0	0.0	0.0	0.0	0.3
<i>Brama japonica</i>	0.0	0.0	0.0	0.0	0.0	1.5	1.5
<i>Pseudopentaceros wheeleri</i>	315.7	1912.4	464.1	13.8	391.4	44.4	3141.8
<i>Pentaceros japonicus</i>	0.0	0.0	0.2	0.0	0.0	0.0	0.2
<i>Antigonia capros</i>	0.0	0.7	4.6	0.0	0.0	0.0	5.4
<i>Antigonia spp.</i>	0.0	0.0	0.2	0.0	0.0	0.0	0.3
<i>Parapercis sp.</i>	0.0	0.5	0.0	0.0	0.0	0.0	0.5
<i>Promethichthys prometheus</i>	47.0	12.8	0.4	1.3	71.3	0.0	132.7
<i>Thysitoides marleyi</i>	0.0	16.3	12.5	0.0	0.0	0.0	28.8
<i>Evoxyemetopon sp.</i>	0.0	1.0	9.8	1.1	0.7	0.0	12.6
<i>Benthodesmus pacificus</i>	0.0	0.0	1.5	0.0	0.0	0.0	1.5
<i>Callionymidae spp.</i>	0.0	0.1	1.3	0.0	0.0	7.1	8.4
<i>Hyperoglyphe japonica</i>	0.0	17.6	7.8	0.0	13.8	0.0	39.2
<i>Microstomus shuntovi</i>	0.0	0.0	2.3	7.2	3.8	53.0	66.2
<i>Chascanopsetta prorigera</i>	0.0	0.4	15.0	0.4	0.9	0.0	16.7
<i>Parabothus coarctatus</i>	0.0	90.2	10.5	0.0	0.0	0.0	100.6
<i>Sphoeroides pachygaster</i>	0.0	0.1	11.8	0.0	0.0	0.0	11.9